

Monthly Water Situation Report August 2018

Natural Resources Wales

- The monthly rainfall total for Wales during August was 100% of the Long Term Average (LTA, 1961-90). South East, South West and North Wales received 106%, 110% and 83% of the LTA, respectively.
- At the end of August, the differences between soil moisture deficit (SMD) values and the LTA across Wales were from -16.7 to 84.7 mm. Soil in 21 squares (out of 23) squares was drier than the LTA for August.
- For river flows in Wales, 17 out of 29 indicator sites (which had flow data available) were classed as *Normal*. 6 sites were *Below normal* and 2 sites were *Notably low*. 3 sites were classed as *Exceptionally low*. The remaining site was *Above Normal*.
- The cumulative reservoir storage for 10 out of 18 indicator reservoirs was greater than 60% at the end of August.

Rainfall*

The monthly rainfall total for Wales was 100% of the LTA for August. The percentage of rainfall recorded in catchments compared with their LTA across Wales was between 51% (Ynys Mon) and 122% (Valleys and Vale of Glamorgan). The rainfall total for Wales was 0.5mm less than the August LTA. For South East, South West and North Wales the rainfall totals were 106%, 110% and 83% of LTA, respectively.

| Geographic regions | Latest Month: July | Last 2 months: June - July | Last 3 months: May - July | Last 6 months: Feb - July | Last 12 months: Aug 2017 - July 2018 | Ranking for current month since 1910 | Ranking for the last two month since 1910 | Ranking for the last three month since 1910 |
|--------------------|--------------------------|-------------------------------------|------------------------------------|------------------------------------|-----------------------------------------------------|-----------------------------------------------------|----------------------------------------------------------|---------------------------------------------------------------|
| | % LTA | | | | | | | |
| North | 83 | 81 | 65 | 69 | 104 | 31 | 22 | 5 |
| South-west | 110 | 99 | 76 | 102 | 109 | 57 | 41 | 13 |
| South-east | 106 | 96 | 72 | 105 | 104 | 49 | 33 | 9 |
| Wales | 100 | 92 | 71 | 98 | 105 | 47 | 32 | 8 |

Rainfall Map <u>National</u>

Rainfall Charts National & Areas South East Wales North Wales South West Wales

All data are provisional and may be subject to revision.

^{*} using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright)

Soil Moisture Deficit/Recharge

The differences between the soil moisture deficits and the LTA for the 23 MORECS squares were from -16.7 to 84.7 mm and soil in 21 squares (out of 23) was drier than the LTA for August.

SMD Map <u>National</u>

SMD Charts Compare to LTA

River Flows

River flows were between *Exceptionally low* and *Above Normal* for all the indicator sites across Wales. 17 out of 29 indicator sites (which had flow data available) were classed as *Normal*. Six sites were *Below normal*, *two* sites were *Notably low*, three sites were classed as *Exceptionally low* and the remaining site was *Above normal* for August.

South East: Flows in the area ranged from 35% (River Lugg at Butts Bridge) to 73% (River Ely at St Fagans) of the August LTA values.

South West: The river flows within this area ranged from 26% (River Teifi at Glanteifi) to 140% (River Llynfi at Coytrahen) of the August LTA values.

North: Flows in the area ranged from 11% (River Cefni at Bodffordd) to 103% (River Conwy at Cwmlanerch) of the August LTA values.

River Flow Map National

River Flow Table % of LTA and compare to previous year

River Flow Charts South East Wales North Wales South West Wales

Groundwater Levels

Groundwater levels for August at indicator sites (9 data available sites) were classed between *Exceptionally low* (Eastwick and Pont y Cambwll) to *Normal* (Pant-y-Lladron, Fernbank, Greenfield Garage, Dodleston Obs and Broxton Obs). The remaining two sites were classed as *Below normal* (Llanfair DC and Handley).

Groundwater Map National

Groundwater Charts South East Wales North Wales South West Wales

Reservoir Storage

At the end of August the the cumulative reservoir storage for 10 out of 18 indicator reservoirs were greater than 60% full.

ReservoirSouth EastNorthSouth WestChartsWalesWalesWales

All data on Water Situation Reports are provisional, based on spot readings, and are subject to revision.

For our latest dry weather update please refer to our webpage here:

 $\underline{\text{https://naturalresources.wales/guidance-and-advice/environmental-topics/water-management-and-quality/dry-weather-update/?lang=en}$

Author: Zhong Zhang Telephone: 03000 654521

All data are provisional and may be subject to revision.

The views expressed in this document are not necessarily those of the Natural Resources Wales. Its officers, servants or agents accept no liability for any loss or damage arising from the interpretation or use the information, or reliance upon views contained herein.

Natural Resources Wales




Figure 1: The Natural Resources Wales Water Situation Report features sites in the catchments shown. Parts of the Shropshire Plains and Mid Severn catchments are within Wales. For full information on these catchments, please see the Environment Agency Midlands Water Situation Report.

For areas adjoining Natural Resources Wales, please see the reports for Environment Agency Midlands and North West England:

<u>Environment Agency - Midlands, England Water Situation Report</u> Environment Agency - North West, England Water Situation Report

All data are provisional and may be subject to revision.

The views expressed in this document are not necessarily those of the Natural Resources Wales. Its officers, servants or agents accept no liability for any loss or damage arising from the interpretation or use the information, or reliance upon views contained herein.

Rainfall





Figure 2: Calculated catchment average August rainfall totals as a percentage of the 1961-90 August long term average for Natural Resources Wales catchments, using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright).

All data are provisional and may be subject to revision.

The views expressed in this document are not necessarily those of the Natural Resources Wales. Its officers, servants or agents accept no liability for any loss or damage arising from the interpretation or use the information, or reliance upon views contained herein.


Rainfall Charts

Figure 3: Rainfall Charts: National and Areas




Comparison of monthly rainfall totals to the 1961-90 long term average expressed as percentage for Natural Resources Wales and Areas, using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright).

Figure 4: Rainfall Charts: South East Wales




Comparison of monthly rainfall totals to the 1961-90 long term average expressed as percentage for South East Wales, using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright).

Figure 5: Rainfall Charts: North Wales




Comparison of monthly rainfall totals to the 1961-90 long term average expressed as percentage for North Wales, using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright).

Return to Summary



Comparison of monthly rainfall totals to the 1961-90 long term average expressed as percentage for North Wales, using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright).

Figure 6: Rainfall Charts: South West Wales



Comparison of monthly rainfall totals to the 1961-90 long term average expressed as percentage for South West Wales, using NCIC (National Climate Information Centre) data (Source: Met Office © Crown Copyright).

Soil Moisture Deficit (SMD)




Figure 7: MORECS soil moisture deficits (mm) for August for real land use for Natural Resources Wales (Source: Met Office © Crown Copyright).

Return to Summary




Figure 8: MORECS month end soil moisture deficits difference (mm) from the 1961-90 long term monthly average (LTA) for August for real land use for Natural Resources Wales squares (Source: Met Office © Crown Copyright).




Figure 9: MORECS month end soil moisture deficit difference (mm) from the 1961-90 long term monthly average (LTA) for real land use for South East, North and South West (Source: Met Office © Crown Copyright). (Note: no LTA available for Natural Resources Wales.)

River Flow




Figure 10: Monthly mean river flow for August, classed relative to analysis of historic August monthly means (Source: Natural Resources Wales).


Return to Summary

| | | August 2018 | | | August 2017 | | August LTA | | |
|------------------------------------|--------------------|-------------------|-------------|----------------|-------------|----------------|------------|--------------------------|--------------------------|
| SITE NAME | RIVER | Class | % of LTA | Flow (m3/s) | % of LTA | Flow (m3/s) | LTA | Monthly Min (m3/s) | Monthly Max (m3/s) |
| River Flow Sites : South East Area | | | | | | | | | |
| Butts Bridge | Lugg | Exceptionally low | 35% | 0.66 | 43% | 0.83 | 1.90 | 0.51 | 6.97 |
| Grosmont | Monnow | Normal | 51% | 0.87 | 72% | 1.22 | 1.69 | 0.36 | 8.17 |
| Pont ar Yscir | Yscir | Normal | 43% | 0.31 | 131% | 0.94 | 0.72 | 0.10 | 3.23 |
| Pontypridd | Taff | Normal | 71% | 7.86 | 131% | 14.50 | 11.10 | 2.29 | 49.00 |
| Redbrook | Wye | Below normal | 49% | 12.60 | 105% | 27.30 | 25.95 | 5.18 | 79.70 |
| Rhiwderin | Ebbw | Below normal | 49% | 1.65 | 118% | 3.95 | 3.35 | 0.93 | 14.30 |
| St Fagans | Ely | Normal | 73% | 1.70 | 109% | 2.54 | 2.34 | 0.46 | 11.30 |
| Trostrey Weir | Usk | Normal | 65% | 5.81 | 146% | 13.10 | 8.96 | 2.70 | 27.50 |
| River Flow Sites : North Area | | | | | | | | | |
| Bodfari | Wheeler | Below normal | 63% | 0.25 | 68% | 0.27 | 0.40 | 0.19 | 0.87 |
| Bodffordd | Cefni | Notably low | 11% | 0.01 | 133% | 0.12 | 0.09 | 0.00 | 0.36 |
| Brynkinalt Weir | Ceiriog | Exceptionally low | 27% | 0.30 | 95% | 1.07 | 1.13 | 0.18 | 5.20 |
| Cwmlanerch | Conwy | Normal | 103% | 11.70 | 152% | 17.40 | 11.41 | 0.73 | 37.40 |
| Cynefail | Gelyn | Normal | 97% | 0.38 | 177% | 0.69 | 0.39 | 0.03 | 1.09 |
| Dol y Bont | Leri | | 0% | 0.00 | | | 1.03 | 0.08 | 3.00 |
| Druid | Alwen | Normal | 90% | 1.90 | 185% | 3.93 | 2.12 | 0.39 | 5.90 |
| Dyfi bridge | Dyfi | | 0% | 0.00 | | | 12.52 | 0.66 | 40.40 |
| Garndolbenmaen | Dwyfor | Normal | 92% | 1.90 | 120% | 2.47 | 2.06 | 0.12 | 6.25 |
| Manley Hall | Dee | Below normal | 67% | 10.20 | 145% | 22.20 | 15.32 | 7.08 | 38.60 |
| Pont y Cambwll | Clwyd | Notably low | 41% | 0.81 | 185% | 3.68 | 1.99 | 0.51 | 7.18 |
| Ruthin Weir | Clwyd | Exceptionally low | 15% | 0.04 | 285% | 0.74 | 0.26 | 0.05 | 0.74 |
| River Flow Sites : South West Area | | | | | | | | | |
| Capel Dewi | Tywi | Normal | 67% | 12.90 | 147% | 28.20 | 19.16 | 2.70 | 78.50 |
| Clog y Fran | Taf | Normal | 66% | 2.34 | 189% | 6.70 | 3.54 | 0.33 | 18.30 |
| Coytrahen | Llynfi | Above normal | 140% | 2.26 | 114% | 1.83 | 1.61 | 0.26 | 6.06 |
| Felin Mynachdy | Cothi | Normal | 81% | 5.05 | 152% | 9.49 | 6.25 | 0.36 | 23.40 |
| Glanteifi | Teifi | Below normal | 26% | 3.46 | 194% | 26.30 | 13.56 | 1.13 | 77.30 |
| Keepers Lodge | Ewenny | Normal | 74% | 0.77 | 98% | 1.02 | 1.04 | 0.22 | 4.24 |
| Marcroft | Afan | Normal | 128% | 4.79 | 142% | 5.34 | 3.75 | 0.55 | 13.30 |
| Pont Llolwyn | Ystwyth | Below normal | 41% | 1.27 | 198% | 6.15 | 3.11 | 0.18 | 8.57 |
| Treffgarne * | Western Cleddau | Normal | 74% | 0.99 | 203% | 2.72 | 1.34 | 0.24 | 5.54 |
| Resolven | Neath | | | | 171% | 9.84 | 5.76 | 0.40 | 22.50 |
| Tir-y-Dail | Loughor | Normal | 82% | 1.00 | 147% | 1.79 | 1.22 | 0.18 | 4.40 |
| Ynystanglws | Tawe | Normal | 112% | 8.71 | 128% | 9.99 | 7.80 | 1.07 | 28.60 |

Figure 11: Monthly mean river flow for August with comparison against previous year expressed as a percentage of the August long term average and classed relative to analysis of historic August monthly means. (Source: Natural Resources Wales). (* For Treffgarne station the LTAs were derived using scaled historical flows (1965-2003) from the downstream station at Prendergast Mill. There was no flow data for Resolven due to the maintainance work at the gauge station)


River Flow Charts

Figure 12: River Flow Charts: South East Wales



Monthly mean river flows for the last 10 months classed relative to the analysis of historic river levels (Source: Natural Resources Wales).

Figure 13: River Flow Charts: North Wales



Monthly mean river flows for the last 10 months classed relative to the analysis of historic river levels (Source: Natural Resources Wales).

(Please note that there was no data for Garndolbenmaen for April to June 2018 due to maintenance work)






Figure 14: River Flow Charts: South West Wales



Return to Summary



Monthly mean river flows for the last 10 months classed relative to the analysis of historic river levels. (Source: Natural Resources Wales).

(* Please note that for Treffgarne station the ranking bands were derived using scaled historical flows (1965-2003) from the downstream station at Prendergast Mill. There were no flow data for June and August 2018 for Resolven)

Groundwater Levels





Figure 15: Groundwater levels at the end of month classed relative to an analysis of historic August groundwater levels (Source: Natural Resources Wales and Environment Agency).


Groundwater charts

Figure 16: Groundwater level charts: South East Wales




End of month groundwater levels for the past 10 months for index sites (Source: Natural Resources Wales). (Please note that data is not available for May and July 2018 for Pant-y-Lladron)

Figure 17: Groundwater level charts: South West Wales



End of month groundwater levels for the past 10 months for index sites (Source: Natural Resources Wales).


Figure 18: Groundwater level charts: North Wales



End of month groundwater levels for the past 10 months for index sites (Source: Natural Resources Wales and Environment Agency). (Please note that data is not available from May to August 2018 for Hollybush)


Reservoir Storage

Figure 19: Reservoir charts: South East Wales




Weekly reservoir stocks for Natural Resources Wales index sites (Source: Welsh Water)

Figure 20: Reservoirs charts: North Wales



Weekly reservoir stocks for Natural Resources Wales index sites (Source: Welsh Water).

Figure 21: Reservoirs charts: South West Wales



Weekly reservoir stocks for Natural Resources Wales index sites (Source: Welsh Water).

Glossary

| Term | Definition | | | | |
|-----------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|--|
| Aquifer Areal average rainfall | A geological formation able to store and transmit water. The estimated average depth of rainfall over a defined area. Expressed in depth of water (mm). The rainfall available to percolate into the soil or produce river flow. Expressed in depth of water (mm). | | | | |
| Effective rainfall | | | | | |
| Groundwater Meteorological Office Rainfall and Evaporation Calculating System (MORECS) | The water found in an aquifer The Met Office provides climate data for grid squares measuring 40km by 40km across the UK using MORECS | | | | |
| Recharge | The process of increasing the water stored in the saturated zone of an aquifer. Expressed in depth of water (mm). | | | | |
| Reservoir live capacity Soil moisture deficit (SMD) | The reservoir capacity normally usable for storage to meet established reservoir operating requirements. It is the total capacity less that not available because of operating agreements or physical restrictions. Only under abnormal conditions, such as a severe water shortage might this additional water be extracted. The difference between the amount of water actually in the | | | | |
| | soil and the amount of water that the soil can hold. Expressed in depth of water (mm). | | | | |
| Categories Exceptionally high Notably high Above normal Normal Below normal Notably low Exceptionally low | Value likely to fall within this band 5% of the time Value likely to fall within this band 8% of the time Value likely to fall within this band 15% of the time Value likely to fall within this band 44% of the time Value likely to fall within this band 15% of the time Value likely to fall within this band 8% of the time Value likely to fall within this band 5% of the time | | | | |
| Units cumecs mAOD | Cubic metres per second (m³ s⁻¹) Metres Above Ordnance Datum (mean sea level at Newlyn Cornwall). | | | | |