

# Know Your River - Glaslyn

## Salmon & Sea Trout Catchment Summary

### Introduction

This report describes the status of the salmon and sea trout populations in the Glaslyn catchment. Bringing together data from rod catches, stock assessments and juvenile monitoring, it will describe the factors limiting the populations and set out the challenges faced in the catchment.


Action tables set out habitat improvements to restore freshwater productivity of salmon and sea trout populations. These tables also include some work which will be carried out by our partner organisations, not just Natural Resources Wales (NRW).

NRW has a duty, defined in the Environment (Wales) Act 2016 to have Sustainable Management of Natural Resources (SMNR) at the core of everything that we do. By applying the principles of SMNR in all of our activities - from agriculture, forestry and flood defence to development planning - we are undertaking catchment-wide initiatives that will deliver for fish stock improvements. Our reports highlight the importance of considering the whole catchment when identifying and addressing fisheries issues; and of working with partners.

NRW is committed to reporting on the status of salmon stocks in all of our principal salmon rivers for the Salmon Action Plans and condition assessments under the Habitats Directive in SAC rivers; all fish species in all of our rivers are reported for the Water Framework Directive (WFD). This report will fulfil these commitments and provide an informative and useful summary of stock status and remedial work planned, for our customers, specifically anglers, fishery and land owners; as well as our partners.

### Catchment

The Glaslyn is predominantly rural with sheep farming in the upper reaches of the catchment, gradually changing to mixed sheep and beef further down the valley. The Glaslyn valley holds a small number of large scale dairy units. There is a large area of afforestation situated on the afon Colwyn.




The soils are glacial in origin and are generally thin, poor, peaty and acidic, but are enriched in places by mineral ores from underlying rocks. The evidence of small scale mineral working of copper and zinc is widespread throughout the Glaslyn catchment, however there are no active metal mines at present. The Sygun copper mine at Beddgelert has been opened as a tourist attraction.

This mountainous area, directly open to the prevailing south westerly winds, receives heavy rainfall. Although the acidity of rainwater is not high, the volume of rain falling in the area gives rise to high annual acid deposition. In the absence of neutralising bases within the thin soils, these areas are 'acid vulnerable' and as a result their water courses are subjected to periodic acid flushes which can reduce water quality and salmonid fish survival. In addition, the abundance of old mine workings, slate quarries and coniferous afforestation are known to exacerbate the impact of acidification in parts of the Glaslyn catchment.


### Rod catches

The following graphs show the total declared rod catches of salmon and sea trout on the Glaslyn.

**Salmon rod catch** – The salmon rod catch in 2017 was still below the ten year average. The release rate was 88%, which is excellent and must continue to conserve stocks.



**Sea trout rod catch** – despite a slight increase compared to 2016, it is still below the ten year average. The release rate in 2017 was 81%. This is excellent and needs to be maintained to improve stocks.




## Stock status

### Conservation of Salmon

Salmon stock status is assessed using 'Conservation Limits' which provide an objective reference point against which to assess the status of salmon stocks in individual rivers.

This is calculated by applying assumed angling exploitation rates to catch data to derive run estimates; adopting standard sex ratios and weight-fecundity relationships to generate egg deposition figures. The numbers of salmon a river can produce (and consequently the catches that the stocks support) are a function of the quality and quantity of accessible spawning and rearing area. Therefore, in general, big rivers have larger catches and have correspondingly bigger total spawning requirements than small rivers. Thus, for any given rivers there should be an optimum level of stock which the conservation limit seeks to protect. The conservation limit represents the number of eggs that must be deposited each year within a given catchment in order to conserve salmon stocks in the future.



Are enough salmon eggs being deposited to conserve salmon stocks in the catchment?

The red line represents the number of eggs required to be deposited to sustain a healthy salmon stock. The black trend line and its confidence limits (the yellow band) is fitted to the most recent 10-year series of egg deposition estimates (2008-2017).


- Current number of eggs being deposited puts stocks **probably at risk**
- In 5 years' time the predicted status of salmon stocks will be **probably at risk**
- Based on current data, and the projection of the graph, the stocks of salmon on the Glaslyn will continue to **decline (uncertain)**

## Conservation of Sea Trout

In contrast to salmon, no established methods of setting Conservation Limits or similar have been available for sea trout. In the absence of such analysis, NRW and the Environment Agency have, for several years, routinely applied a fishery based assessment to the principal sea trout rivers. This method – used previously in this report - utilises time-series' of angling catch per unit effort (CPUE) data ('catch per day') to examine sea trout performance on a river-by-river basis.

Recently an alternative stock-based assessment method has been developed by NRW and is applied here. This utilises angling catch data to derive run and egg deposition estimates for sea trout in much the same way that similar data sets are used in Conservation Limit compliance procedures for salmon assessment.

Further details on this method are given in the recent Technical Case supporting net and rod fishery byelaw proposals on all rivers in Wales and the cross-border rivers Wye and Dee (see: <http://naturalresourceswales.gov.uk/media/682258/technical-case-structure-final.pdf>)



Are enough sea trout eggs being deposited to conserve stocks in the catchment?

The red line represents the number of eggs required to be deposited to sustain a healthy sea trout stock. The black trend line and its confidence limits (the yellow band) is fitted to the most recent 10-year series of egg deposition estimates (2008-2017).

- Current number of eggs being deposited puts stocks **probably at risk**
- In 5 years' time the predicted status of salmon stocks will be **probably at risk**
- Based on current data, and the projection of the graph, sea trout stocks will continue to **decline (uncertain)** on the Glaslyn

### **Juvenile Monitoring**

The monitoring season was hindered in 2017 by wet weather and high flows. This led to the temporal site on the Dwyfor not being completed. This was also the case for most temporal sites in Gwynedd and Meirionnydd.